

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 July 2005 (07.07.2005)

PCT

(10) International Publication Number
WO 2005/061442 A1

(51) International Patent Classification⁷: **C07C 235/46**,
C07D 211/46, 213/82, 401/12, 401/14, 405/12, 409/12

(21) International Application Number:
PCT/US2004/038227

(22) International Filing Date: 6 December 2004 (06.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/529,061 12 December 2003 (12.12.2003) US

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, Indiana 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BENESH, Dana, Rae** [US/US]; 13287 Beckwith Drive, Westfield, Indiana 46074 (US). **BLANCO-PILLADO, Maria-Jesus** [ES/US]; 7278 North Hawthorne Lane, Indianapolis, Indiana 46250 (US).

(74) Agents: **GINAH, Francis, O.** et al.; Patent Division, Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: OPIOID RECEPTOR ANTAGONISTS



(57) Abstract: A compound of the formula (I) wherein the variables X_1 to X_5 , R^1 to R^7 including R^3 , E, q, v, y, z, A and B are as described, or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixtures thereof, useful for the treatment, prevention or amelioration of obesity and Related Diseases is disclosed.



WO 2005/061442 A1



— *of inventorship (Rule 4.17(iv)) for US only*

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

OPIOID RECEPTOR ANTAGONISTS

The present invention is in the field of medicinal chemistry. The invention relates specifically to compounds useful as opioid antagonists, methods of treatment, methods of using, and pharmaceutical compositions thereof.


Background

Three types of opioid receptors, mu, kappa, and delta opioid receptors are generally reported. Recent evidence points to the interactions between receptor dimer combinations of mu, kappa and/or delta receptors (called heterodimers) as also contributing to opioid activity. Opioid receptors and their normal regulation or lack thereof, has been implicated in disease states including irritable bowel syndrome, nausea, vomiting, pruritic dermatoses, depression, smoking and alcohol addiction, sexual dysfunction, stroke and trauma in animals. Therefore it is not surprising that the ability to antagonistically bind opioid receptors has been shown to produce ameliorative, preventative and/or treatment effects in animals including humans afflicted with one or more of these disease states.


More recently, certain antagonists of the opioid receptors have been found to increase metabolic energy consumption, and reduction of weight in obese rats while maintaining muscle mass. These findings indicate that an effective opioid antagonist may be useful in preventing, treating and/or ameliorating the effect of obesity. Considering the percentage of the population that is obese in Western societies and the indirect costs associated with treating the effects and symptoms of obesity and Related Diseases, the importance of these findings cannot be overstated.

Though many opioid antagonists have been disclosed, the search continues for alternative and/or improved or more effective antagonists having an overall benefit to the patient with little or no major side effects. U.S Patent No. 4,891,379 disclosed phenylpiperidine opioid antagonists useful for the treatment of diabetes and obesity. In particular, U.S. patent 4,891,379 disclosed the compound LY 255582 represented by the structure:

2




U.S. Patent No. 4,191,771 also disclosed compounds useful as opioid antagonists. Also, bicyclic analogs of phenyl piperidine have been prepared and reported as opioid antagonists in Wentland, et al., *Biorganic and Medicinal Chemistry Letters* 11 (2001) 623-626; see also Wentland, et al., *Bioorganic and Medicinal Chemistry Letters* 11 (2001) 1717-1721. Finally, European Patent application number EP 1 072592A2 filed May 18, 2000, discloses phenylpiperidine compounds of formula 1



1

wherein A, D, R¹, R², R³, X, and n have meanings given in the description, which are useful in the prophylaxis and in the treatment of diseases mediated by opioid receptors such as pruritus.

U.S patent No. 6,140,352 and related patents disclose the compound of formula Formula 1




(1)

wherein the variables X_1 , X_2 , X_3 , R_1 , R_3 , R_4 , R_5 and R_6 are as described therein, as agonists of the beta adrenergic receptor useful for the treatment of diabetes and obesity.

Regardless of these and other disclosures of compounds useful as opioid receptor antagonists, or useful for the treatment of obesity, and/or diabetes by other mechanisms, there remains an unmet medical need for a safe, effective and/or alternate treatment or prophylaxis of diseases associated with opioid receptors, particularly obesity and Related Diseases.

Summary of the Invention

The present invention provides a compound of the formula I



wherein

each of X_1 , X_2 , X_3 , X_4 , and X_5 is C, CH, or N; provided that ring B has no more than 2 nitrogen atoms;

X is NH or CH_2 , so that ring A is cyclohexyl, cyclohexenyl, or piperidinyl;

E is NH or O;

v is 0, 1, 2, or 3;

q is 0 or 1, provided that when the A-ring is cyclohexyl or cyclohexenyl q is 1 and provided that v and q are not simultaneously 0;

R^1 and R^2 are independently selected from hydrogen, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, aryl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_{10}$ alkylaryl, heterocyclyl, $\text{C}_1\text{-C}_{10}$ alkylheterocyclic, $\text{-C}_1\text{-C}_8$ alkylC(O) $\text{C}_1\text{-C}_8$ alkyl, $\text{-(CH}_2)_n\text{(CO) C}_3\text{-C}_8$ cycloalkyl-, $\text{-C}_2\text{-C}_8$ alkylCH(OH)aryl, -, $\text{-CO(O)C}_1\text{-C}_8$ alkyl, $\text{-SO}_2\text{C}_1\text{-C}_8$ alkyl, $\text{-SO}_2\text{C}_1\text{-C}_{10}$ alkylaryl, $\text{-SO}_2\text{C}_1\text{-C}_8$ alkylheterocyclic, $\text{-C}_1\text{-C}_8$ alkylcycloalkyl, $\text{-(CH}_2)_n\text{C(O)OR}^8$, $\text{-(CH}_2)_n\text{C(O)R}^8$, $\text{-(CH}_2)_m\text{C(O)NR}^8\text{R}^8$, and $\text{(CH}_2)_m\text{NSO}_2\text{R}^8$; wherein each of the alkyl, alkenyl, cycloalkyl, heterocyclic, and aryl groups are optionally substituted with one to five groups independently selected from halo, $\text{C}_1\text{-C}_8$ haloalkyl, $\text{C}_1\text{-C}_8$ thioalkyl, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, aryl, $\text{-C}_1\text{-C}_8$ alkylaryl,

-C(O)C₁-C₈ alkyl, -SO₂C₁-C₈ alkyl, -SO₂C₁-C₈ alkylaryl, -C₁-C₈ alkylcycloalkyl; and wherein R¹ and R² may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of amino, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, aryl, C₁-C₈ alkylaryl, -C(O)C₁-C₈ alkyl, -CO(O)C₁-C₈ alkyl, halo, oxo, C₁-C₈ haloalkyl;

R³ and R^{3'} are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, aryl, -C₁-C₈ alkylcycloalkyl, or -C₁-C₈ alkylaryl; C₁-C₈ alkylheterocyclic; or R³ and R^{3'} combine to form a C₃-C₈ cycloalkyl, C₄-C₈ cycloalkenyl, or C₅-C₁₀ heterocyclic;

R⁴ and R⁵ are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, -C₂-C₈ alkynyl, -C₁-C₈ alkoxyalkyl, C₁-C₈ thioalkyl, halo, C₁-C₈ haloalkyl, -C₁-C₈ alkoxyhaloalkyl, aryl, -C₁-C₈ alkylaryl, -C(O)C₁-C₈ alkyl, or -C(O)OC₁-C₈ alkyl, -C₁-C₈ alkylamino, -C₁-C₈ alkylcycloalkyl, -(CH₂)_mC(O)C₁-C₈ alkyl, and (CH₂)_nNR⁸R⁸, wherein each R⁴ or R⁵ is attached to its respective ring only at carbon atoms, and wherein y is 0, 1, 2, or 3; and wherein z is 0, 1, 2, or 3;

R⁶ and R⁷ are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -C(O)C₁-C₈ alkyl, hydroxy, C₁-C₈ alkoxy, -SO₂C₁-C₈ alkyl, SO₂C₁-C₈ alkylaryl, -SO₂C₁-C₈ alkylheterocyclic, aryl, -C₁-C₈ alkylaryl, C₃-C₇ cycloalkyl, -C₁-C₆ alkylcycloalkyl, -(CH₂)_nC(O)R⁸, -(CH₂)_mC(O)NR⁸R⁸, and -(CH₂)_mNSO₂R⁸; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C₁-C₈ alkyl, C₂-C₈ alkenyl, aryl, and C₁-C₈ alkylaryl; and wherein R⁶ and R⁷ may independently combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may optionally have substituents selected from the group consisting of oxo, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, aryl, -C₁-C₈ alkylaryl, -C(O)C₁-C₈ alkyl, -CO(O)C₁-C₈ alkyl, hydroxy, C₁-C₈ alkoxy, -C₁-C₈ alkylamine, amino, halo, and haloalkyl;

R⁸ is hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₁-C₈ alkylaryl, -C(O)C₁-C₈ alkyl, or -C(O)OC₁-C₈ alkyl; and wherein n is 0, 1, 2, 3 or 4 and m is 1, 2, or 3;

or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

The present invention also provides a pharmaceutical formulation comprising a compound of formula I in association with a carrier, diluent and/or excipient.

The present invention also relates to a method for the treatment and/or prophylaxis of obesity and Related Diseases including eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, depression related to obesity, anxiety related to obesity, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, stroke, metabolic diseases and symptoms thereof, hyperlipidemia, hypertriglycemia, hyperglycemia, hyperlipoproteinemia, substance abuse, drug overdose, compulsive behavior disorders (such as paw licking in dog), and addictive behaviors such as for example, gambling, and alcoholism, comprising administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

The present invention provides a compound of formula I useful for the manufacture of a medicament for the treatment, prevention and/or amelioration of symptoms associated with obesity and Related Diseases.

In another embodiment, the present invention provides a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixtures thereof, useful as an appetite suppressant.

In another embodiment, the present invention provides a method of achieving weight loss while maintaining or minimizing the loss of lean muscle mass, comprising administering a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixtures thereof, to a patient in need thereof.

In yet another embodiment, the present invention provides a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture thereof, to a patient in need thereof in combination with other effective therapy for the treatment of obesity and related disorders.

Detailed Description of the Invention

As used herein the term "obesity" has its commonly understood meaning such as "excessively fat" and includes the clinical designation of being obese as defined in and by the medical literature and brochures of support or public health organizations. For

example, *Dorland's Illustrated Medical Dictionary* (29th edition, W.B. Saunders Company, Philadelphia USA.) defines obesity as "an increase in bodyweight beyond the limitation of skeletal and physical requirements, as the result of an excessive accumulation of fat in the body." Because the decision of suitability administration of a compound(s) of the present invention to a patient is to be made by a qualified physician or qualified caregiver, the patient is inherently deemed suitable or obese by the administering caregiver.

As used herein, the term "patient" includes human and non-human animals such as companion animals (dogs and cats) and livestock animals.

The preferred patient of treatment, amelioration and/or prevention of obesity and Related Diseases are human.

The terms "treating" and "treat", as used herein, include their generally accepted meanings, *i.e.*, preventing, prohibiting, restraining, alleviating, ameliorating, slowing, stopping, or reversing the progression or severity of a pathological condition, or sequela thereof, described herein.

The terms "ameliorating" "preventing", "prevention of", "prophylaxis", "prophylactic" and "prevent" are used herein interchangeably and refer to reducing the severity of the symptoms associated with obesity and Related Diseases in a patient afflicted with same or reducing the likelihood that the recipient of a compound of formula I will incur or develop any of the pathological conditions, or sequela thereof, described herein.

As used herein, the term "effective amount" is synonymous with "effective dose" and means an amount of a compound of formula I that is sufficient in one or more administrations for preventing, ameliorating or treating a condition, or detrimental effect thereof, herein described, or an amount of a compound of formula I that is sufficient for antagonizing the opioid receptors to achieve the desired outcome within the purview of the invention.

The term "pharmaceutically acceptable" is used herein as an adjective and means substantially non-deleterious to the recipient patient.

The term "Active Ingredient" as used herein means a compound of formula I or a combination of compounds of formula I or a combination of a compound of formula I and

a co-antagonist of the opioid receptor or a combination of a compound of formula I in addition to other effective anti-obesity, weight loss or anti-diabetic agent(s).

The term "formulation", as in pharmaceutical formulation, or "pharmaceutical composition" is intended to encompass a product comprising the Active Ingredient (as defined supra), and the inert ingredient(s) that make up the carrier, or other components of the drug as administered, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the Active ingredients, or from dissociation of one or more of the Active ingredients. Accordingly, the pharmaceutical formulations of the present invention encompass any effective composition made by admixing a compound of the present invention and a pharmaceutical carrier. The pharmaceutical formulations of the present invention also encompass a compound of the formula I and a pharmaceutically acceptable co-antagonist of opioid receptors or other effective therapy useful for the treatment and/or prevention of obesity or Related Diseases.

The term "Related Diseases" as used herein refers to such symptoms, diseases or conditions caused by, exacerbated by, induced by or adjunct to the condition of being obese. Such diseases, conditions and/or symptoms include but are not limited to eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, obesity related depression, obesity related anxiety, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, stroke, metabolic diseases and symptoms thereof, hyperlipidemia, hypertriglycemia, hyperglycemia, and hyperlipoproteinemia. As used herein the terms obesity related depression and obesity related anxiety are conditions of depression and anxiety respectively, that are symptomatic of certain obese patients and possibly brought on by the awareness or self-consciousness of the condition of being obese and possibly coupled with the real or perceived notion of acceptance or rejection by the certain individual, individuals or the public at large. Obesity related depression or anxiety may generally be alleviated or treated adjunctively with the underlying condition of being obese or overweight and/or prevented by administration of a compound of formula I.

The term "suitable solvent" refers to any solvent, or mixture of solvents, inert to the ongoing reaction that sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction.

The term "mutual solvent" means a solvent that is used to dissolve sufficiently, two or more components of a reaction or mixture separately prior to reaction or mixing, that is a solvent common to more than one reagents or components of a mixture.

The term "nitrogen containing heterocycle" refers to a aromatic or non-aromatic, monocyclic or bicyclic ring system which is a 4, 5, 6, or 7-member ring containing 1, 2 or 3 nitrogen atoms in addition to the carbon atoms completing the ring size, or a combination of 1 nitrogen atom and 1, or 2 atoms selected from oxygen, and sulfur in addition to the appropriate number of carbon atoms completing the ring size. A nitrogen containing heterocycle as used here may have 0, 1, 2 or 3 double bonds.

The term "C₁-C₈ alkyl" or C₁₋₈ alkyl" refers to and includes all groups, structural isomers and /or homologues of alkyl groups having from 1 to 8 carbon atoms. When the term C₁-C₈ alkyl precedes or prefixes another group, the term C₁-C₈ alkyl, only limits the number of carbon atoms in the alkyl component. For example C₁-C₈ alkyaryl means an aryl group having a C₁-C₈ alkyl group substituent such that the number of carbon atoms in the group C₁-C₈ alkylaryl is effectively the number of carbon atoms in the aryl group plus the number of carbon atoms in the C₁-C₈ alkyl group. Similarly, the term "C₁-C₈ alkylcycloalkyl" refers to a cycloalkane group having a C₁-C₈ alkyl substituent, and wherein the entire group C₁-C₈ alkylcycloalkane may itself be a substituent attached at either the alkyl group or the cycloalkyl group to a substrate. The definition and usage applies equally to other homologues of C₁-C₈ such as for example, C₁-C₇, C₁-C₆ etc. In general, and where necessary a dash (-) has been placed next to certain groups to indicate the point of attachment for clarity. Nevertheless, the absence of a dash does not otherwise negate the position obvious position(s) of attachment known to one of skill in the art.

The term "cycloalkane" or "cycloalkyl" means cycloalkanes having from 3 to 8 carbon atoms i.e. from cyclopropane to cyclooctane.

The term "hal" or "halo" as used herein refers to a halogen including fluorine, chlorine, bromine or iodine.

The term "haloalkane" or "haloalkyl" means haloalkanes having from 1 to 8 carbon atoms, and from 1 to 3 halogen atoms as allowed by valency considerations. Examples include chloroethyl, trifluoromethyl, 2-chloropropyl, etc.

As used herein the terms "alkenyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon double bonds.

As used herein the terms "alkynyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon triple bonds.

As used herein the term "alkoxy" refers to the group "O-alkyl" wherein alkyl is as defined previously.

The term "aryl" as used herein refers to compounds or groups having the Huckel $4n+2$ pi electron arrangement and includes for example, phenyl, benzyl, naphthyl, tetrahydronaphthyl, benzothiophene, etc, but excludes carbazoles and other fused tricyclic ring structures.

As used herein the term "aroxy" or "aryloxy" refers to the group "O-aryl" wherein aryl is as defined previously.

As used herein the term "fused bicyclic" means a fused cycloalkane ring system wherein each ring has from 4 to 8 carbon atoms (i.e. C_8 - C_{16} fused bicyclic) and the fused ring system has from 0 to 3 bridgehead carbon atoms. One or both of the fused rings may contain zero or one double bond. Examples of fused bicyclics include but are not limited to bicyclo[2,2,1]heptyl, bicyclo[2,2,1]heptenyl.

As used herein the term "heterocyclic" or "heterocyclyl" or "heterocycle" are used interchangeably and has its usual meaning and includes mono, bi or tricyclic or spirocyclic heterocyclic groups unless otherwise specified. Heterocycles as used herein may contain 1, 2, or 3 heteroatoms selected independently from nitrogen, oxygen or sulfur, unless otherwise specified. Examples of heterocyclic groups applicable to the present invention include but are not limited to pyranyl, piperazinyl, pyrrolidinyl, azapanyl, azaflorenyl, isoquinolinyl, indolinyl, thiophenyl, benzothiophenyl, oxazolyl, morpholinyl, thiomorpholinyl, and piperidinyl. Each of the heterocyclic groups may be mono or di substituted or as specified with substituents such as alkyl, cycloalkyl, aryl, among others as defined. Furthermore, substitution may be at the 1-position or heteroatom as in piperazine, pyrrolidine or at a carbon atom or both.

As used herein, the term "protecting group" refers to a group useful for masking reactive sites in a molecule to enhance the reactivity of another group or allow reaction at another desired site or sites following which the protecting group may be removed. Protecting groups are usually used to protect or mask groups including but not limited to -OH, -NH, and -COOH. Suitable protecting groups are known to one of skill in the art and are described in *Protecting groups in Organic Synthesis*, 3rd edition, Greene, T. W.; Wuts, P.G.M. Eds., John Wiley and Sons, New York, 1999.

As used herein, the term "solvate" is a form of the compound of the invention wherein a crystal or crystals of a compound of the invention have been formed from a stoichiometric or non-stoichiometric amount of the compound of formula I and a solvent. Typical solvating solvents include for example, water, methanol, ethanol, acetone and dimethylformamide.


In those instances where a compound of the invention possesses acidic or basic functional groups, various salts may be formed which are more water soluble and/or more physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, calcium, magnesium, aluminum and the like. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion-exchange resin.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," *J. Phar. Sci.*, 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, hydrobromide, camsylate, carbonate, clavulanate, citrate, chloride, edetate, edisylate, estolate, esylate, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrochloride, hydroxynaphthoate, hydroiodide, isothionate, lactate, lactobionate, laurate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pantothenate, phosphate, polygalacturonate,

salicylate, stearate, subacetate, succinate, tannate, tartrate, tosylate, trifluoroacetate, trifluoromethane sulfonate, and valerate. Preferred salts for the purpose of the invention include the hydrochloride salt, the hydrobromide salt, the bisulfate salt, the methane sulfonic acid salt, the *p*-toluenesulfonic acid salt, bitartrate, the acetate and the citrate salt.

A compound of the invention as illustrated by formula I may occur as any one of its positional isomers, stereochemical isomers or regio- isomers, all of which are objects of the invention. Certain compounds of the invention may possess one or more chiral centers, and thus, may exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group, there exist the possibility of *cis*- and *trans*- isomeric forms of the compounds. The *R*- and *S*- isomers and mixtures thereof, including racemic mixtures as well as mixtures of enantiomers or *cis*- and *trans*- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art by using stereospecific reactions with starting materials which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods. For example, a racemic mixture may be reacted with a single enantiomer of some other compound i.e. a chiral resolving agent. This changes the racemic form into a mixture of stereoisomers and diastereomers, because they have different melting points, different boiling points, and different solubilities and can be separated by conventional means, such as crystallization.

PCT international application WO 02/078693 A2 published October 10, 2002 discloses compounds of the formula



wherein R₁, R₂, R₃, R₄ and X are as described therein, as antagonists of the 5-HT₆ receptor for the treatment of disorders including cognitive disorders, age related disorders, mood disorders, psychosis, etc. The compounds of the present invention however, are useful for the treatment and/or prevention of obesity and Related Diseases. The compounds of the present invention have also shown inhibition of orexigenic effects,


and are thus useful as appetite suppressants either as a single therapy or as combination therapy in conjunction with exercise and other effective appetite suppressing or weight loss medications.

The efficacy of certain compounds of the present invention have been demonstrated by their activity or potency in several biological models including a binding scintillation proximity assay (SPA) and functional GTP-gamma-S assay).


Preferred Embodiments of the Invention

A compound of formula I preferably exists as the free base or a pharmaceutically acceptable salt. More preferred is the hydrochloride salt, the bisulfate salt, mesylate or the oxalic acid salt of the compound of formula I.


Preferred embodiments of the compound of formula I include the substructures Ia, Ib Ic and Id as shown below:




(Ia);



(Ib);




(Ic);



For the groups R^1 and R^2


Preferred R^1 and R^2 groups are independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, phenyl, naphthyl, benzothiophene, and isopropyl.

Also preferred are R^1 and R^2 groups independently selected from the group consisting of methyl, ethyl, propyl, isopropyl, phenyl,



each of which is optionally substituted with a group selected from the group consisting of halogen, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 thioalkyl, C_1 - C_8 alkylamino, phenyl, C_1 - C_8 alkylsubstituted phenyl, C_4 - C_8 heterocycle or $-C_1$ - C_4 alkylheterocycle; or combine with a group selected from C_1 - C_8 alkyl, halogen, C_1 - C_8 haloalkyl, C_1 - C_8 thioalkyl, C_1 - C_8 alkylamino, phenyl, C_1 - C_8 alkylsubstituted phenyl, C_4 - C_8 heterocycle or C_1 - C_4 alkyl heterocycle to form a substituted or unsubstituted bicycle or tricycle, and wherein n is preferably 1, 2, or 3.

Also preferred are R^1 and R^2 groups that combine with each other or with 1 or 2 atoms adjacent to the nitrogen atom to form a group selected from the group consisting of



each of which is optionally substituted with a group selected from the group consisting of halogen, amino, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, -C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or -C₁-C₄ alkylheterocycle.

Preferred R³ and R^{3'} Groups

A preferred R³ is hydrogen. A preferred R^{3'} group is selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, phenyl and benzyl.

Preferred R⁴ Groups

A preferred R⁴ group is selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ haloalkyl, C₁-C₅ alkoxy, -C₁-C₅ alkylamino, -N(C₁-C₅ alkyl)₂, -NHC₁-C₅ alkyl, -C₁-C₅ alkyl N(C₁-C₅ alkyl)₂, -C₁-C₅ alkylNHC₁-C₅ alkyl, phenyl, -C₁-C₅ alkylphenyl, -C₁-C₅ alkylcycloalkyl, and C₁-C₅ thioalkyl. More preferred is a R⁴ group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. Most preferred is an R⁴ group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, propoxy, isopropoxy, and benzyl.

Though the groups R⁴ and a R⁵ may exist as multiple substituents on their respective ring substrates, a preferred embodiment of the invention involves compounds wherein each of R⁴, and R⁵ are independently singly or doubly substituted on their respective ring substrates.

Preferred R⁵ Groups

A preferred R⁵ group is selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ haloalkyl, C₁-C₅ alkoxy, -C₁-C₅ alkylamino, -N(C₁-C₅ alkyl)₂, -NHC₁-C₅

alkyl, -C₁-C₅ alkylN(C₁-C₅ alkyl)₂, -C₁-C₅ alkylNHC₁-C₅ alkyl, phenyl, -C₁-C₅ alkylphenyl, -C₁-C₅ alkylcycloalkyl, and C₁-C₅ thioalkyl. More preferred is an R⁵ group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. A most preferred R⁵ group is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, and benzyl.

Preferred R⁶ and R⁷ Groups

Preferred are R⁶ and R⁷ groups independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, isopropyl, phenyl and benzyl.

Also preferred are compounds of formula I wherein R⁶ and R⁷ independently combine with each other, and with the nitrogen atom to which they are attached or with 1, or 2 atoms adjacent to the nitrogen atom to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle may optionally have substituents selected from the group consisting of oxo, amino, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, -C₁-C₈ alkylaryl, -C(O)C₁-C₈ alkyl, -CO(O)C₁-C₈ alkyl, hydroxy, C₁-C₈ alkoxy, halo, and haloalkyl.

Preferred values for q

q is preferably 0 when the A-ring is piperidinyl. P is preferably 1 when the A-ring is cyclohexyl.

Preferred E group

A most preferred E group is an oxygen atom (O).

Preferred A-ring

A preferred A-ring is a cyclohexyl or piperidinyl. Most preferred A ring is cyclohexyl.

Preferred B-ring

A preferred B-ring is a phenyl ring, a pyrazine ring, a pyrimidine ring or a pyridine ring. Most preferred B ring is a phenyl, pyrazine or pyridine ring.

Preferred values for v, n and m


A preferred value for v is 0, 1, or 2.

A preferred value for n is 1, 2 or 3.


A preferred value for m is 1 or 2.

A preferred compound according to the present invention is a compound selected from the group consisting of:


\pm 6-[4-[2-(tetrahydro-pyran-4-yl)-ethylamino]-cyclohexyloxy]-nicotinamide,




\pm 6-[4-(3-Methyl-butylamino)-cyclohexyloxy]-nicotinamide,




\pm 6-[4-(2-Thiophen-2-yl-ethylamino)-cyclohexyloxy]-nicotinamide




\pm 4-[4-(3-Phenyl-propylamino)-cyclohexyloxy]-benzamide




Trans-6-(4-Benzylamino-cyclohexyloxy)-nicotinamide,




6-(1-Pyridin-2-ylmethyl-piperidin-4-yloxy)-nicotinamide




6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-nicotinamide




6-[1-(1H-Indol-2-ylmethyl)-piperidin-4-yloxy]-nicotinamide



4-(1-Benzyl-piperidin-4-yloxy)-benzamide,



4-[1-(3-Phenyl-propyl)-piperidin-4-yloxy]-benzamide




and a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer or a diastereomeric mixture thereof.

Preparing Compounds of the Invention

A typical protocol for the preparation of compounds of the invention and intermediates thereof wherein the A-ring is an optionally substituted cyclohexyl group is depicted in Scheme 1 below.

Scheme 1




According to Scheme 1, the starting material 3,3-Dimethyl-1,5-dioxaspiro[5.5]undecanone (**1**) and analogs thereof is reduced to the corresponding alcohol 3,3-Dimethyl-1,5-dioxaspiro[5.5]undecan-9-ol (**2**) by reaction with polymer supported borohydride in a methanolic solvent. One of skill in the art is aware that other reducing agents and modes of reduction (e.g. without polymer support) may be utilized to afford the compound **2** and analogs thereof. The ketone **1** and analogs thereof may be purchased from Chemical distributors such as for example, Aldrich Chemical Co, Milwaukee, USA. The compound 3,3-Dimethyl-1,5-dioxaspiro[5.5]undecan-9-ol (**2**) is then coupled with halo nicotinonitrile or halobenzonitrile or other B-ring source to afford the oxygen linked compound **3** or analog thereof. For example, optionally substituted 4-chloronicotinonitrile is reacted with compound **2** to afford the oxygen-linked compound **3** under basic conditions. Basic conditions include the use of bases selected from inorganic and organic bases. Examples of useful inorganic bases include but are not limited to potassium carbonate, sodium hydride, sodium carbonate, sodium hydroxide, potassium hydroxide, calcium carbonate and cesium carbonate. Examples of organic bases include but are not limited to potassium hexamethyl disilazide, n-butyl lithium, hexamethylphosphorous triamide, (HMPT), and the like. The basic conditions are complemented by the presence of a solvent, preferably an organic solvent. Preferred organic solvents include protic solvents or polar aprotic solvents. Most preferred solvents include dimethylformamide, methanol, dimethylacetamide (DMA), dimethylsulfoxide. A most preferred basic reaction condition involves the use of potassium carbonate in dimethylacetamide at temperatures of about 60 to 100 °C. The protecting group (dimethylacetal group) of compound **3** is removed by reaction with an acidic group such as for example, hydrochloric acid to afford the compound **4**. The compound **4** is reductively aminated with a desired amine to afford the amino compound **5**, which is a compound of the invention. The reductive amination may be performed in two steps or a single step depending on the stability of the intermediate imine intermediate. Typically, compound **4** is reacted with a primary or secondary amine (primary amine shown) in methanol as solvent. Molecular sieves may be added to enhance the efficiency of the imine formation. In a second step the reducing agent, typically, sodium borohydride or other hydride reducing agent is added to the reaction mixture. The progress of the reaction may be monitored by TLC, HPLC, LC-MS or other analytical technique known

to one of skill in the art to determine the substantial completion of each step and timing for the addition of the next reagent. The resulting amino nitrile compound **5** is hydrolyzed at the cyano group to afford the primary amide **6**. Nitrile hydrolysis is preferably accomplished by reaction with hydrogen peroxide and an inorganic base such as sodium carbonate and preferably under pressure. A suitable solvent for accomplishing the above nitrile hydrolysis is DMSO or DMF.

Analogues of compounds **3** and **5** having one or more substituent R groups may be prepared by using appropriately substituted starting materials or by inter-conversion of substituent functionality. For example an initial optional substituent group on the A or B ring may be protected and deprotected appropriately to achieve the desired end substituent R. Alternatively an initial substituent may be converted by known 1, 2 or 3 step reactions to other desired final substituents.

An alternate protocol illustrated in Scheme 2 shows the use of the benzamide as the source of ring B.

Scheme 2




The use of the amide starting material is particularly preferred for compounds of the invention where the B-ring is pyridinyl, pyridazinyl, pyrazinyl or pyrimidinyl group.

The carboxamide or heterocyclic amide may be introduced as part of the starting material where the appropriate surrogate for the B-ring is commercially available or may be prepared using known methods. For example, the use of pyrazine carboxamide, nicotinamide or substituted analogs thereof results in substituted derivatives or analogs of compounds of formula **3a** or **6a**, which are also compounds of the present invention.

Primary and secondary amines are useful for the reductive amination to convert compound **4a** to compound **6a** as shown in Scheme 2. Examples of useful amines for the reductive amination include but are not limited to phenethylamine, 3-methylbutylamine, propylamine, isopropylamine, benzylamine and isopentylamine.

Compounds prepared by this and other schemes disclosed herein, or known to one of skill in the art may further be converted to the acid addition salt as shown for example, in Scheme 3.

Scheme 3




Scheme 3 shows preparation of the hydrochloride salt **12a**, a compound of the invention wherein R¹R²NH is 3-methylbutylamine or other secondary amine group and R⁴ and R⁵ are both hydrogen. As shown, the starting material **7** is 4-hydroxypiperidine

protected at the nitrogen atom using tertiary butoxycarbonyl anhydride (Boc-anhydride). The Boc-protected piperidinol (**7**) is reacted with a source of the B-ring such as a halobenzonitrile, or a haloniconitrile (6-chloro-nicotinonitrile (**8**) shown) or halopyridazino nitrile or carboxamide thereof as desired. The coupling reaction to form the ether linkage (**9**) is performed in the presence of a base such as sodium hydride or sodium carbonate in a suitable solvent such as DMA, DMF, or DMSO. The nitrile group of the resulting ether (**9**) is then hydrolyzed to form the primary amide. Hydrolysis of the nitrile is accomplished in the presence of hydrogen peroxide and a base such as sodium carbonate. The resulting amide **10** is hydrolyzed under acidic conditions to afford the deprotected compound **11**. Deprotection of the Boc group is best accomplished using HCl, TFA or HF. Procedures for Boc-protection and deprotection are known to one of skill in the art and are described in general Organic chemistry references including *Protecting groups in Organic Synthesis*, 3rd edition, Greene, T. W.; Wuts, P.G.M. Eds., John Wiley and Sons, New York, 1999. Specific procedures may also be found in the experimental section herein. The free piperidinyl NH group of compound **11** may be reacted with an aldehyde having the desired alkyl, alkylaryl, cycloalkyl, alkylcycloalkyl, alkylheterocyclic or other substituent within the scope of the invention to afford the desired N-substituted piperidinyl compound **12**.

The compound **12** is dissolved in ethanol and a slight excess (e.g 1.0 to 1.5 molar equivalents based on the number of basic sites) of 1N hydrochloric acid is added at temperatures ranging from about 0 °C to room temperature. The mixture may be allowed to crystallize over time with or without cooling, or may be evaporated to afford the hydrochloride salt, which may be further purified by trituration with a suitable organic solvent such as toluene, hexanes, diethylether or mixtures thereof. Alternatively, anhydrous HCl may be bubbled into a cold solution of compound **12** until the reaction is complete or the solution is saturated, and the mixture worked up as appropriate to afford compound **12a**. One of skill in the art is aware of the nuances and the varied techniques for preparing, isolating and purifying acid addition salts, and should achieve comparable results using methods appropriate for the particular substrate without undue experimentation.

A modified protocol for preparing compounds of the invention is provided in Scheme 4 wherein the nucleophilic displacement reaction to form the ether linkage is performed towards the end of the synthesis rather than early on.

Scheme 4




According to Scheme 4, the starting material is an appropriately substituted hydroxypiperidine (7) protected at the nitrogen using Boc anhydride. It may be possible to purchase the Boc-protected hydroxy piperidine. The Boc protected hydroxypiperidine 7 is reacted with a B-ring source such as 4-fluorobenzonitrile to afford the ether linked compound 9. Other B ring sources include for example, phenyl or pyridine carboxamide, benzonitrile or pyridino-nitrile and analogs thereof. Methods of accomplishing the coupling reaction have been disclosed previously. The compound 9 may be hydrolyzed

to the amide via the nitrile group, deprotected by removing the Boc group as disclosed previously, and finally reductively aminated to afford a compound of the invention. Alternatively, the ether compound **9** may be deprotected by removing the Boc group to afford the compound **13**. The protecting group may be removed by use of hydrochloric acid or trifluoroacetic acid using procedures known to one of skill in the art. One of skill in the art is aware that appropriately substituted analogs of the compound of formula **13** may be prepared by starting with appropriately substituted starting materials or surrogates thereof which may be converted to the desired substituents.

Deprotection of compound **9** to form compound **13** is followed by reductive amination to form the N-substituted piperidinyl compound **14**. The N-substituted piperidinyl compound **14** is finally hydrolyzed at the nitrile group to afford compound **15**, a compound of the invention.


Compounds of formula I wherein v is 1 may be made following the synthetic scheme described below:



As shown in Scheme 5 4-hydroxy-cyclohexanecarboxylic acid ethyl ester (commercially available from Aldrich Chemical Company, Milwaukee, USA or other fine chemical suppliers) may be reacted with a source of the B-ring such as halobenzonitrile or

haloniconitrile to form the ether linked product **13**. The coupling reaction to form the ether linkage is performed in the presence of a base such as sodium hydride or potassium carbonate in a suitable solvent such as DMA, DMF or DMSO. The carboxylic acid ester is then selectively reduced to give the corresponding aldehyde **14**. This reduction is accomplished with hydrides such as for example diisobutylaluminum hydride (DIBAL-H). The aldehyde **14** is then reductively aminated with the desired amino moiety to form the amine **15**. The nitrile of the resulting amino precursor is then hydrolyzed to yield a compound of the invention **16**.

Yet another protocol for the preparation of compounds of formula I is shown in Scheme 6.



The aldehyde **14** of scheme 5 is reacted with methoxymethyldiphenyl phosphine oxide or methoxymethyltriphenylphosphonium chloride in the presence of a strong base such as *n*-butyl lithium, *sec*-butyl lithium or potassium hexamethyldisilane or the like to afford the vinyl methylether **17**. The vinylmethyl ether **17** is then hydrolyzed under acidic conditions to afford the higher aldehyde **18**. The aldehyde **18** is then converted to the desired compound **20** of formula I as shown and discussed previously.

Method of Using the Invention

As noted above, the compounds of the present invention are useful in blocking the effect of agonists at mu, kappa, and/or delta opioid receptors. As such, the present invention also provides a method for blocking a mu, kappa, delta receptor or receptor combination (heterodimer) thereof in a mammal comprising administering to said mammal a receptor blocking dose of a compound of formula I.

The term "receptor blocking dose", as used herein, means an amount of a compound of formula I necessary to block a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof following administration to a mammal requiring blocking of a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof.

The compounds of formula I or combinations thereof, are effective over a wide dosage range. For example, dosages per day will normally fall within the range of about 0.05 to about 250 mg/kg of body weight. In the treatment of adult humans, the range of about 0.5 to about 100 mg/kg, in single or divided doses, is preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician in light of the relevant circumstances, including the condition to be treated, the choice of compound to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration. Therefore, the above dosage ranges are not intended to limit the scope of the invention in any way. The compounds may be administered by a variety of routes such as the oral, transdermal, subcutaneous, sublingual, intranasal, intramuscular and intravenous routes.

A variety of physiologic functions have been shown to be subject to or influenced by mu, kappa, or delta receptors or receptor combination (heterodimers) in the brain. As such, the compounds of the present invention are believed to have the ability to treat disorders associated with these receptors or combinations thereof, such as eating disorders, opioid overdose, depression, smoking, alcoholism, sexual dysfunction, shock, stroke, spinal damage and head trauma. As such, the present invention also provides methods of treating the above disorders by blocking the effect of agonists at a mu, kappa, delta receptors or receptor combinations (heterodimer) thereof. The compounds of the present invention have been found to display excellent activity in an opioid receptor

binding assay which measures the ability of the compounds to block the mu, kappa, delta or receptor combination (heterodimer) thereof.

GTP- γ -S Binding Assay

An SPA - based GTP- γ -S assay format was developed based on previous opioid (Emmerson et al., J. Pharm Exp Ther 278,1121,1996; Horng et al., Society for Neuroscience Abstracts, 434.6, 2000) and muscarinic (DeLapp et al., JPET 289, 946, 1999) assay formats. Membranes were re-suspended in 20 mM HEPES, 100 mM NaCl, 5 mM MgCl₂, 1 mM DTT, and 1 mM EDTA. Fifty (50) mL of GTP- γ -[35S], compound, membrane suspension (20 microgram/well), and wheat germ agglutinin coated SPA beads (1mg/well) were added to clear bottom 96 well assay plates. GDP (200 mM) was added to the membrane solution prior to addition to the assay plates. Plates were sealed and incubated for four hours at room temperature then placed in a refrigerator overnight to allow the beads to settle. Signal stability at 4 °C was determined to be > 60 hours. Plates were warmed to room temperature and counted in a Wallac Microbeta scintillation counter. For antagonist assays, specific agonists were added at the following concentrations: (MOR) DAMGO 1 micromolar, (DOR) DPDPE 30 nM, (KOR) U69593 300 nM. Kb's were determined by Cheng-Prusoff equation (see Cheng and Prusoff, Biochem. Pharmacol. 22, 3099, 1973). Results obtained for a representative sample of compounds of the invention in the GTP- γ -S Binding Assay are shown in table 1 below.

Table 1

In Vitro Antagonism GTP- γ -S Binding Assay

<u>Compound of</u>	<u>K_b (uM)</u>	<u>K_b (uM)</u>	<u>K_b (uM)</u>
<u>Example #</u>	<u>Mu</u>	<u>Kappa</u>	<u>Delta</u>
1	0.566	0.601	0.819
2	0.594	0.620	0.996
6	>1.800	0.430	>7.300
7	1.785	1.051	>7.300
8	>1.850	0.937	>7.300

9	1.120	0.930	>7.300
10	0.453	0.999	1.965

Ex-Vivo Receptor Binding

In order to bridge in vitro binding affinity and antagonist potency to in vivo potency and efficacy applicants have developed an ex vivo receptor binding assay in rat brain. This assay measures the difference in association (binding) of a high affinity nonselective opioid receptor radioligand (3H-diprenorphine) in brain tissue isolated from animals receiving vehicle versus compound treatment (less binding of 3H-diprenorphine = greater compound association with opioid receptors). Studies using the ex-vivo receptor binding assay have demonstrated a positive correlation between activity (potency and duration of activity) which also correlates to 24 hour efficacy in dietary induced obese rats.

Methods. An opioid receptor ex vivo binding assay measures 3H-diprenorphine binding (0.1 –0.4 nM affinity radioligand for mu, delta and kappa receptors) in rat striatum/nucleus accumbens; a region of the brain that contains a high density of mu, delta and kappa receptors, following oral administration of compounds. Experimentally, a screening dose of 7 mg/kg, p.o. of compound or vehicle is administered to rats. Six hours following compound administration, the animals are sacrificed and the striatum/nucleus accumbens is isolated and homogenized in 10 volumes (weight/volume) binding buffer. The homogenate is then used in a homogenate binding assay using a saturating concentration of 3H-diprenorphine for 30 minutes. The homogenization and assay is performed at 4 °C, to minimize compound redistribution in the in vitro binding portion of the assay. Results are reported (Table 2) as specific binding constant K_i in micromolar (μ M).

Table 2
SPA Binding Affinity assay K_i (μ M)

Compound of Example No.	Mu	Kappa	Delta
3	0.137	2.561	0.353

4	0.620	>5.000	2.345
---	-------	--------	-------

Formulation

A compound of the invention is preferably presented in the form of a pharmaceutical formulation comprising a pharmaceutically acceptable carrier, diluent or excipient and a compound of the invention. Such compositions will contain from about 0.1 percent by weight to about 90.0 percent by weight of the compound of the invention (Active Ingredient). As such, the present invention also provides pharmaceutical formulations comprising a compound of the invention and a pharmaceutically acceptable carrier, diluent or excipient thereof.

In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material that acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, emulsions, solutions, syrups, suspensions, aerosols (as a solid or in a liquid medium), and soft and hard gelatin capsules.

Examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, water, and mineral oil. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

For oral administration, the Active Ingredient, a compound of this invention, may be admixed with carriers and diluents and molded into tablets or enclosed in gelatin capsules.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 1 to about 500 mg, more usually about 5 to about 300 mg, of the Active Ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

In order to more fully illustrate the operation of this invention, the following formulation examples are provided. The examples are illustrative only, and are not intended to limit the scope of the invention. The formulations may employ as Active Ingredient any of the compounds of the present invention.

FORMULATION 1

Hard gelatin capsules are prepared using the following ingredients:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	250	55
Starch dried	200	43
Magnesium stearate	10	2

The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities.

FORMULATION 2

Capsules each containing 20 mg of medicament are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	20	10
Starch	89	44.5
Microcrystalline cellulose	89	44.5
Magnesium stearate	2	1

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve and filled into a hard gelatin capsule.

FORMULATION 3

Capsules each containing 100 mg of active ingredient are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	100	30
Polyoxyethylene Sorbitan monooleate	50mcg	0.02
Starch powder	250	69.98

The above ingredients are thoroughly mixed and placed in an empty gelatin capsule.

FORMULATION 4

Tablets each containing 10 mg of active ingredient are prepared as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	10	10
Starch	45	45
Microcrystalline cellulose	35	35
Polyvinylpyrrolidone (as 10% solution in water)	4	4
Sodium carboxymethyl starch	4.5	4.5
Magnesium stearate	0.5	0.5
talc	1	1

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granule so produced is dried at 50-60 °C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules, which after mixing, is compressed on a tablet machine to yield a tablet weighing 100 mg.

FORMULATION 5

A tablet formula may be prepared using the ingredients below:

Compound	Amount per capsule (mg)	Percent by weight (%)
Active Ingredient	250	38
Cellulose microcrystalline	400	60
Silicon dioxide fumed	10	1.5
Stearic acid	5	0.5

The components are blended and compressed to form tablets each weighing 665mg.

FORMULATION 6

Suspensions each containing 5 mg of medicament per 5 mL dose are made as follows:

Compound	Amount per 5mL suspension (ml)
Active Ingredient	5
Sodium carboxymethyl cellulose	50
Syrup	1.25
Benzoic acid solution	0.10
Flavor	q.v.
Color	q.v.
Water	q.s. to 5mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethylcellulose and syrup to form a smooth paste. The benzoic acid

solution, flavor and color is diluted with some of the water and added to the paste with stirring. Sufficient water is then added to produce the required volume.

FORMULATION 7

An aerosol solution is prepared containing the following components:


Compound	Concentration by weight (percent)
Active Ingredient	0.25
Ethanol	29.75
Propellant 22 (chlorodifluoromethane)	70.0

The active compound is mixed with ethanol and the mixture added to a portion of the Propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted further with the remaining amount of propellant. The valve units are then fitted to the container.

Examples


Example 1

\pm 6-[4-[2-(Tetrahydro-pyran-4-yl)-ethylamino]-cyclohexyloxy]-nicotinamide,



Step 1


Preparation of 3,3-Dimethyl-1,5-dioxaspiro[5.5]undecan-9-ol



Combine 3,3-dimethyl-1,5-dioxaspiro[5.5]undecan-9-one (Aldrich, 750 mg, 3.78 mmol) with polymer-supported borohydride (Aldrich, 3026 mg, 7.56 mmol) in methanol (30 mL). Shake by rotation the resulting mixture overnight. Filter the reaction mixture and concentrate the filtrate. Wash the residue with hydrochloric acid 0.1M and extract with EtOAc. Dry the organic layer over sodium sulfate, filter and concentrate.

Step 2


Preparation of 6-(3,3-Dimethyl-1,5-dioxaspiro[5.5]undec-9-yloxy)-nicotinonitrile



Add dropwise a solution of 3,3-dimethyl-1,5-dioxaspiro[5.5]undecan-9-ol, (1445 mg, 7.22 mmol) in DMF (2.1 mL) to a suspension of sodium hydride (433 mg, 10.82 mmol) in DMF (8.6 mL). Let the reaction mixture stir at room temperature for 1h, then heat while stirring at 50°C for 20 min. Add dropwise a solution of 6-chloro-nicotinonitrile (1200 mg, 8.66 mmol) in DMF (4.5 mL). Continue the heating at 60°C and stirring overnight. Concentrate the reaction mixture to remove DMF. Wash the residue with water (15 mL) and extract with EtOAc/hexanes (20 mL). Dry the organic layer over sodium sulfate, filter and concentrate. Purify the residue by flash chromatography (eluent CH₂Cl₂/hexanes 2/1) to give 2000 mg (92% yield) of the title compound.

Step 3


Preparation of 6-(4-Oxo-cyclohexyloxy)-nicotinonitrile



Combine hydrochloric acid (1.0M aq., 20 mL) with a solution of 6-(3,3-dimethyl-1,5-dioxaspiro[5.5]undec-9-yloxy)-nicotinonitrile (2000 mg, 6.61 mmol) in acetone (25 mL). Stir at room temperature for 2h then at 40-50°C for 1h. Concentrate the reaction mixture. Partition the residue between EtOAc/hex (25 mL) and K₂CO₃ (aq. sat. 20 mL). Wash the organic layer with water, brine, and dry it over sodium sulfate, filter and concentrate. Triturate the residue with EtOAc/hexanes (1/4) to provide a white solid which is further purified by flash chromatography (EtOAc/Hexanes 1/4) to give 1010 mg (71% yield) as a white solid.

Step 4

Preparation of 6-{4-[2-(Tetrahydro-pyran-4-yl)-ethylamino]-cyclohexyloxy}-nicotinonitrile, NE4-A05445-035




Combine the previously obtained 6-(4-oxo-cyclohexyloxy)-nicotinonitrile (, 200 mg, 0.925 mmol), 2-(tetrahydro-pyran-4-yl)-ethylamine (Aldrich, 125 mg, 0.971 mmol) and a scoop of molecules sieves 3Å in methanol (4 mL). Let the reaction mixture stir overnight, and then add sodium borohydride (70 mg, 1.85 mmol). Continue the stirring overnight. Purify the reaction mixture by loading onto an SCX column, washing with methanol and eluting with ammonia/methanol (2.0M). Purify the residue by two flash chromatographies (40/1 CH₂Cl₂/ammonia in methanol) to provide 199 mg (65%) of the title compound as pale yellow oil.

Step 5

Combine the previously obtained 6-{4-[2-(tetrahydro-pyran-4-yl)-ethylamino]-cyclohexyloxy}-nicotinonitrile (199 mg, 0.604 mmol) and potassium carbonate in DMSO (4 mL). Cool the reaction mixture to 0°C, and then add hydrogen peroxide (0.181 mL) dropwise. Let the reaction mixture stir at room temperature for 3h. Pour the mixture onto water. Filter the precipitate and redissolved in methanol. Concentrate and purify through an SCX column to provide the title compound. Mass spectrum (ion spray): $m/z = 348.3$ (M+1); ^1H NMR (CDCl_3): 8.62 (s, 1H), 8.04 (m, 1H), 6.75 (m, 1H), 6.01 (bs, 2H), 5.32 (bs, 1H), 5.07 (m, 1H), 3.97 (dd, $J=11.0$ and 4.0 Hz, 2H), 3.40 (t, $J=11.4$ Hz, 2H), 2.73-2.54 (m, 3H), 2.21-2.03 (m, 3H), 1.79-1.28 (m, 12H).

Example 2


\pm 6-[4-(3-Methyl-butylamino)-cyclohexyloxy]-nicotinamide



Using a method similar to example 1, gives the title compound (199 mg, 96%). Mass spectrum (ion spray): $m/z = 306.3$ (M+1); ^1H NMR (CDCl_3): 8.66 (s, 1H), 8.13-8.10 (m, 1H), 7.96 (bs, 1H), 7.39 (bs, 1H), 6.81 (m, 1H), 5.17 (s, 1H), 5.00 (m, 1H), 2.49 (m, 3H), 2.06-1.14 (m, 14H), 0.88 (t, $J=6.6$ Hz, 6H).

Example 3

\pm 6-[4-(2-Thiophen-2-yl-ethylamino)-cyclohexyloxy]-nicotinamide




Using a method similar to example 1, gives the title compound (203 mg, 87%). Mass spectrum (ion spray): $m/z = 346.1$ (M+1); ^1H NMR (CDCl_3): 8.61 (s, 1H), 8.05-

8.00 (m, 1H), 7.17 (m, 1H), 6.96 (m, 1H), 6.87 (s, 1H), 6.98-6.95 (m, 1H), 6.03 (bs, 2H), 5.31-5.04 (m, 1H), 3.07-2.96 (m, 4H), 2.67-2.55 (m, 1H), 2.20-2.00 (m, 3H), 1.78-1.26 (m, 6H).


Example 4

4-[4-(3-Phenyl-propylamino)-cyclohexyloxy]-benzamide



Step 1

4-(3,3-Dimethyl-1,5-dioxaspiro[5.5]undec-9-yloxy)-benzonitrile




Add dropwise a solution of 3,3-dimethyl-1,5-dioxaspiro[5.5]undecan-9-ol, (NE4-A05445-029, 1377 mg, 6.88 mmol) in DMF (2.0 mL) to a suspension of sodium hydride (412 mg, 10.32 mmol) in DMF (8.0 mL). Let the reaction mixture stir at room temperature for 1h, then heat while stirring at 50°C for 20 minutes (min). Add dropwise a solution of 4-fluoro-benzonitrile (1000 mg, 8.26 mmol) in DMF (4.2 mL). Continue the heating at 60°C and stirring for 2 hours (h). Concentrate the reaction mixture to remove DMF. Wash the residue with water (15 mL) and extract with EtOAc/hexanes (20 mL). Dry the organic layer over sodium sulfate, filter and concentrate. Purify the residue by flash chromatography (eluent CH₂Cl₂/hexanes 2/1) to give xxx mg (xx% yield) of the title compound.

38

Step 2


4-(4-Oxo-cyclohexyloxy)-benzonitrile



Combine hydrochloric acid (1.0M aq., 20 mL) with a solution of in acetone (25 mL). Stir at room temperature for 2 hours then at 40-50°C for 1h. Concentrate the reaction mixture. Partition the residue between EtOAc/hex (25 mL) and K₂CO₃ (aq. sat. 20 mL). Wash the organic layer with water, brine. Dry the organic layer over sodium sulfate, filter and concentrate. Triturate the residue with EtOAc/hexanes (1/4) to provide a white solid which is further purified by flash chromatography (EtOAc/Hexanes 1/4) to give mg (% yield) as a white solid.

Step 3

4-(4-Oxo-cyclohexyloxy)-benzamide



Combine 4-(4-oxo-cyclohexyloxy)-benzonitrile previously obtained (100 mg, 0.464 mmol), K₂CO₃ (32 mg, 0.232 mmol) in DMSO (23 mL). Cool the reaction mixture to 0°C and add hydrogen peroxide (0.139 mL). Let stir the reaction mixture for 4h at room temperature. Quench the reaction mixture with water (15 mL). Extract with EtOAc/hex 2/1 (3x20 mL). Dried over sodium sulfate, filter and concentrate. Purify the residue by flash chromatography (20/1 CH₂Cl₂/ammonia in methanol 2.0M) to provide the title compound (40 mg, 37%).


Step 4

Combine 4-(4-oxo-cyclohexyloxy)-benzamide (20 mg, 0.085 mmol), 3-phenyl-propylamine (11 mg, 0.085 mmol), triacetoxyborohydride (23 mg, 0.111 mmol) and

acetic acid (5 μ L, 0.085 mmol) in CH_2Cl_2 (1 mL). Let stir overnight. Purify by SCX column (ammonia in methanol 2.0M). Triturate the residue with EtOAc/hexanes 1/1 to provide a white powder (25 mg, 86%).


Example 5

Trans-6-(4-Benzylamino-cyclohexyloxy)-nicotinamide



Step 1


4-Benzylamino-cyclohexanol



Combine *trans*-4-amino-cyclohexanol (2.0 g, 17.4 mmol) in methanol (75 mL) in a sealed tube then add benzaldehyde (1.85 mL, 18.23 mmol). Heat the reaction mixture at 70°C while stirring for 2h. Then let the reaction cool down and add sodium borohydride (2.46 g, 65.1 mmol) in portions. Stir overnight. Evaporate the solvent till 1/3 of the original volume. Partition the reaction mixture between EtOAc (50 mL) and water (40 mL). Reextract the aqueous layer with EtOAc (20 mL). Combine the organic layers and dry over sodium sulfate. Filter and concentrate to provide the title compound (3.5 g) that will be use directly in the next step.

Step 2

6-(4-Benzylamino-cyclohexyloxy)-nicotinonitrile



Add dropwise a solution of 4-benzylamino-cyclohexanol (675 mg, 3.28 mmol) in DMF (2 mL) to a suspension of NaH (196 mg, 4.92 mmol) in DMF (3 mL). Stir at room temperature for 45 min then at 50°C for additional 40 min. Add a solution of 6-chloro-nicotinonitrile (500 mg, 3.61 mmol) in DMF (1.8 mL) dropwise and stir overnight at 60°C. Cool down the reaction mixture and evaporate the solvent. Wash the residue with water (10) and extract with EtOAc/hex (2/1, 15 mL). Combine the organic layers and dry over sodium sulfate. Filter and concentrate. Purify the resulting residue through an SCX column. Further purified by chromatography [$\text{CH}_2\text{Cl}_2/\text{NH}_3$ (2.0M in methanol) 20/1] to provide the title compound (890 mg, 88%).


Step 3

Add K_2CO_3 (200 mg, 1.44 mmol) to a solution of 6-(4-benzylamino-cyclohexyloxy)-nicotinonitrile (890 mg, 2.89 mmol) in DMSO (25 mL). Cool the reaction to 0°C and add hydrogen peroxide (0.87 mL). Stir the resulting reaction mixture at room temperature for 2h. Then quench the reaction mixture with water (25 mL) and extract with EtOAc (30 mL). Dried over sodium sulfate, filter and concentrate. Further purify the residue by SCX chromatography to provide the title compound (700 mg, 74%). Mass spectrum (ion spray): $m/z = 326.0$ ($M+1$); ^1H NMR (CDCl_3): 8.61 (s, 1H), 8.04 (m, 1H), 7.36 (m, 5H), 6.74 (d, $J=8.8$ Hz, 1H), 5.10 (m, 1H), 3.87 (s, 2H), 2.64 (m, 1H), 2.22-2.07 (m, 4H), 1.58-1.33 (m, 4H).

Example 6


6-(1-Pyridin-2-ylmethyl-piperidin-4-yloxy)-nicotinamide

41



Step 1


4-(5-Cyano-pyridin-2-yloxy)-piperidine-1-carboxylic acid tert-butyl ester



Add dropwise a solution of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester (1210 mg, 6.01 mmol) in DMF (1.8 mL) to a suspension of NaH (360 mg, 9.02 mmol) in DMF (7.2 mL). Stir at room temperature for 45 min then at 50°C for additional 40 min. Add a solution of 6-chloro-nicotinonitrile (1000 mg, 7.22 mmol) in DMF (3.6 mL) dropwise and stir overnight at 60°C. Cool down the reaction mixture and evaporate the solvent. Wash the residue with water (10) and extract with EtOAc/hex (2/1, 15 mL). Combine the organic layers and dry over sodium sulfate. Filter and concentrate. Purify the resulting residue through an SCX column. Further purified by chromatography [$\text{CH}_2\text{Cl}_2/\text{NH}_3$ (2.0M in methanol) 20/1] to provide the title compound (1.73 g, 94%).

Step 2

4-(5-Carbamoyl-pyridin-2-yloxy)-piperidine-1-carboxylic acid tert-butyl ester




Combine a solution of 4-(5-cyano-pyridin-2-yloxy)-piperidine-1-carboxylic acid tert-butyl ester (1630 mg, 5.38 mmol) in DMSO (50 mL) with potassium carbonate (371

mg, 2.69 mmol). Cool the solution to 0°C and add slowly hydrogen peroxide (1.61 mL). After 10 min, stir the reaction mixture at room temperature for 2h. Add water (25 mL) and extract twice with CH₂Cl₂ (30 mL). Dry the organic layer over sodium sulfate, filter and concentrate to provide the title compound (1669 mg, 97%) as a white solid.

Step 3

6-(Piperidin-4-yloxy)-nicotinamide hydrochloride




Combine 4-(5-carbamoyl-pyridin-2-yloxy)-piperidine-1-carboxylic acid tert-butyl ester (1559 mg, 4.85 mmol) in tetrahydrofuran (25 mL) with hydrochloric acid (4.0 M in dioxane, 15 mL). Stir the resulting reaction mixture for 48h. Filter the white precipitate washing with EtOAc (10 mL). Redissolve the white solid in methanol and concentrate to provide the title compound (1195 mg, 89%).

Step 4

Combine 6-(piperidin-4-yloxy)-nicotinamide (100 mg, 0.45 mmol) with sodium triacetoxo-borohydride (124 mg, 0.59 mmol) and pyridine-2-carbaldehyde (43 µL, 0.45 mmol) in CH₂Cl₂ (1.5 mL). Stir the reaction mixture for 3h. Then, dilute the reaction mixture with CH₂Cl₂ (5 mL) and washed with NaOH (1M aq, 5 mL). Dry the organic layer over sodium sulfate, filter and concentrate. Purify the residue through an SCX chromatography to provide the title compound (83 mg, 59%). Mass spectrum (ion spray): $m/z = 313.1$ (M+1); ¹H NMR (DMSO-*d*₆): 8.67 (ad, J=2.2 Hz, 1H), 8.50 (m, 1H), 8.12 (dd, J=2.6 and 8.8 Hz, 1H), 7.98 (bs, 1H), 7.80-7.76 (m, 1H), 7.47 (d, J=7.9 Hz, 1H), 7.41 (bs, 1H), 7.29-7.26 (m, 1H), 6.84 (d, J=8.8 Hz, 1H), 5.10 (m, 1H), 3.63 (m, 2H), 2.77-2.74 (m, 2H), 2.35-2.30 (m, 2H), 1.99 (bs, 2H), 1.75-1.69 (m, 2H).

Example 7


6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-nicotinamide



Using a method similar to example 6, gives the title compound (89 mg, 72%).
Mass spectrum (ion spray): $m/z = 276.1$ ($M+1$); ^1H NMR ($\text{DMSO}-d_6$): 8.66 (m, 1H), 8.12 (app. dd, $J=2.2$ and 8.3 Hz, 1H), 7.97 (bs, 1H), 7.40 (bs, 1H), 6.84 (app. d, $J=8.3\text{Hz}$, 1H), 5.08-5.03 (m, 1H), 2.82 (bs, 2H), 2.52 (s, 1H), 2.27-2.20 (m, 3H), 1.99 (m, 2H), 1.72 (m, 2H), 0.86-0.83 (m, 1H), 0.50-0.45 (m, 2H), 0.10-0.07 (m, 2H).

Example 8


6-[1-(1H-Indol-2-ylmethyl)-piperidin-4-yloxy]-nicotinamide




Using a method similar to example 6, gives the title compound (110 mg, 70%).
Mass spectrum (ion spray): $m/z = 351.1$ ($M+1$).

Example 9

4-(1-Benzyl-piperidin-4-yloxy)-benzamide

**Step 1**

4-(4-Cyano-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester




Add a solution of N-Boc-4-hydroxypiperidine (3.0 g, 14.9 mmol) in DMF (5 mL) to a suspension of sodium hydride (894 mg, 22.4 mmol) in DMF (17 mL). Stir the reaction mixture while heating at 50°C for 45 min. Then add a solution of 4-fluorobenzonitrile (2.16 g, 17.9 mmol) in DMF (5 mL). Stir and heat at 50°C for 2h. Let cool to room temperature and quench with water (0.5 mL). Evaporate DMF. Redissolved the resulting residue in EtOAc/hexanes (2/1, 20 mL) and wash with water (3x15 mL). Dry the organic layer over magnesium sulfate, filter and concentrate. Purify by chromatography (EtOAc/hexanes 20% and EtOAc/hexanes 10%) to yield the title compound (2.32 g, 52%).

45

Step 2


4-(Piperidin-4-yloxy)-benzonitrile hydrochloride



Add dropwise acetyl chloride (2.5 mL) to methanol (5.0 mL) at 0°C. Stir the resulting solution at 0°C for 90 min. Then add a solution of 4-(4-cyano-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester (284 mg, 0.94 mmol) in methanol. Stir the resulting mixture for 3h. Evaporate the solvent and triturate with diethyl ether to provide the title compound (216 mg, 96%).

Step 3

4-(1-Benzyl-piperidin-4-yloxy)-benzonitrile



Combine 4-(piperidin-4-yloxy)-benzonitrile hydrochloride (64 mg, 0.268 mmol), benzaldehyde (55 μ L, 0.536 mmol) and sodium triacetoxy borohydride (85 mg, 0.402 mmol) in CH_2Cl_2 (3 mL). Stir at room temperature overnight. Dilute the reaction mixture with CH_2Cl_2 (3 mL) and wash with NaOH (1M aq. 5 mL). Separate the organic layer and place it in an SCX column, eluting with ammonia (2.0M in methanol) to provide the title compound (74 mg, 95%).


Step 4

Combine 4-(1-benzyl-piperidin-4-yloxy)-benzonitrile (74 mg, 0.25 mmol), DMSO (2.5 mL) and powdered potassium carbonate (18 mg, 0.13 mmol). Cool the resulting mixture to 0°C and add hydrogen peroxide (76 μ L). After addition, stir the mixture at room temperature for 1h. Quench the reaction mixture with water (2 mL). Filter the precipitate formed rinsing with diethyl ether to give the title compound (57 mg, 73%).

Mass spectrum (ion spray): $m/z = 311.1$ ($M+1$); ^1H NMR (CDCl_3): 7.79 (ad, $J = 8.6$ Hz, 3H), 7.32-7.21 (m, 5H), 7.14 (bs, 1H), 6.95 (d, $J = 8.6$ Hz, 2H), 4.49-4.42 (m, 1H), 3.31 (s, 2H), 2.69-2.62 (m, 2H), 2.27-2.19 (m, 2H), 1.96-1.89 (m, 2H), 1.66-1.56 (m, 2H).


Example 10

4-[1-(3-Phenyl-propyl)-piperidin-4-yloxy]-benzamide



Step 1

4-(4-Carbamoyl-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester




Combine 4-(4-cyano-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester Example 9, step 1, 215 mg, 0.71 mmol) and potassium carbonate (49 mg, 0.36 mmol) in DMSO (35 mL) at 0°C . Then add dropwise hydrogen peroxide (213 μL). Then stir at room temperature for 1h. Quench the reaction with water (10 mL) and extract with EtOAc/hexanes (2/1, 3x20 mL). Combine the organic layers, dry over magnesium sulfate, filter and concentrate to give the title compound.

47

Step 2

4-(Piperidin-4-yloxy)-benzamide hydrochloride




Add hydrochloric acid (4M in dioxane, 3 mL) dropwise to a solution of 4-(4-carbamoyl-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester (228 mg, 0.711 mmol) in THF (2 mL). Stir at room temperature for 7 hours (h). Filter the solids and dry under reduced pressure to give the title compound (180 mg, 99%) as hydrochloride.

Step 3

Combine 4-(piperidin-4-yloxy)-benzamide hydrochloride (102 mg, 0.397 mmol), 3-phenyl-propionaldehyde (106 μ L, 0.796 mmol) and triacetoxy-borohydride (127 mg, 0.597 mmol) in CH_2Cl_2 (3 mL). Stir the reaction mixture at room temperature overnight. Dilute the reaction mixture with CH_2Cl_2 (3 mL) and wash with sodium hydroxide (1N, aq. 5 mL). Separate the organic layer and placed onto an SCX column directly eluting with ammonia (2.0 in methanol). The resulting residue was triturated with CH_2Cl_2 and diethyl ether to provide the title compound (63 mg, 47%) as a white solid. mass spectrum (ion spray): $m/z = 339.1$ ($M+1$); ^1H NMR (CDCl_3): 7.79 (ad, $J = 9.0$ Hz, 3H), 7.28-7.12 (m, 6H), 6.95 (d, $J = 9.0$ Hz, 2H), 4.74-4.40 (m, 1H), 2.70-2.63 (bm, 2H), 2.56 (t, $J = 7.6$ Hz, 2H), 2.31-2.25 (bm, 2H), 2.22-2.15 (bm, 2H), 1.96-1.89 (bm, 2H), 1.71 (pentet, $J = 7.3$ Hz, 2H), 1.63-1.56 (m, 2H).

We claim:

1. A compound of formula (I)



wherein

each of X_1 , X_2 , X_3 , X_4 , and X_5 is C, CH, or N; provided that ring B has no more than 2 nitrogen atoms;

X is NH or CH_2 , so that ring A is cyclohexyl, cyclohexenyl, or piperidinyl;

E is NH or O;

v is 0, 1, 2, or 3;

q is 0 or 1, provided that when the A-ring is cyclohexyl or cyclohexenyl q is 1 and provided that v and q are not simultaneously 0;

R^1 and R^2 are independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, aryl, C_3 - C_8 cycloalkyl, C_1 - C_{10} alkylaryl, heterocyclyl, C_1 - C_{10} alkylheterocyclic, - C_1 - C_8 alkylC(O) C_1 - C_8 alkyl, $-(CH_2)_n(CO) C_3$ - C_8 cycloalkyl-, $-C_2$ - C_8 alkylCH(OH)aryl, -, $-CO(O)C_1$ - C_8 alkyl, $-SO_2C_1$ - C_8 alkyl, $-SO_2C_1$ - C_{10} alkylaryl, $-SO_2C_1$ - C_8 alkylheterocyclic, - C_1 - C_8 alkylcycloalkyl, $-(CH_2)_nC(O)OR^8$, $-(CH_2)_nC(O)R^8$, $-(CH_2)_mC(O)NR^8R^8$, and $-(CH_2)_mNSO_2R^8$; wherein each of the alkyl, alkenyl, cycloalkyl, heterocyclic, and aryl groups are optionally substituted with one to five groups independently selected from halo, C_1 - C_8 haloalkyl, C_1 - C_8 thioalkyl, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, aryl, $-C_1$ - C_8 alkylaryl, $-C(O)C_1$ - C_8 alkyl, $-SO_2C_1$ - C_8 alkyl, $-SO_2C_1$ - C_8 alkylaryl, $-C_1$ - C_8 alkylcycloalkyl; and wherein R^1 and R^2 may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of amino, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, aryl, C_1 - C_8 alkylaryl, $-C(O)C_1$ - C_8 alkyl, $-CO(O)C_1$ - C_8 alkyl, halo, oxo, C_1 - C_8 haloalkyl;

R^3 and $R^{3'}$ are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, aryl, $-C_1$ - C_8 alkylcycloalkyl, or $-C_1$ - C_8 alkylaryl; C_1 - C_8 alkylheterocyclic; or R^3 and $R^{3'}$ combine to form a C_3 - C_8 cycloalkyl, C_4 - C_8 cycloalkenyl, or C_5 - C_{10} heterocyclic;

R^4 and R^5 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-C_1$ - C_8 alkoxyalkyl, C_1 - C_8 thioalkyl, halo, C_1 - C_8 haloalkyl, $-C_1$ - C_8 alkoxyhaloalkyl, aryl, $-C_1$ - C_8 alkylaryl, $-C(O)C_1$ - C_8 alkyl, or $-C(O)OC_1$ - C_8 alkyl, $-C_1$ - C_8 alkylamino, $-C_1$ - C_8 alkylcycloalkyl, $-(CH_2)_mC(O)C_1$ - C_8 alkyl, and $(CH_2)_nNR^8R^8$, wherein each R^4 or R^5 is attached to its respective ring only at carbon atoms, and wherein y is 0, 1, 2, or 3; and wherein z is 0, 1, 2, or 3;


R^6 and R^7 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-C(O)C_1$ - C_8 alkyl, hydroxy, C_1 - C_8 alkoxy, $-SO_2C_1$ - C_8 alkyl, SO_2C_1 - C_8 alkylaryl, $-SO_2C_1$ - C_8 alkylheterocyclic, aryl, $-C_1$ - C_8 alkylaryl, C_3 - C_7 cycloalkyl, $-C_1$ - C_6 alkylcycloalkyl, $-(CH_2)_nC(O)R^8$, $-(CH_2)_mC(O)NR^8R^8$, and $-(CH_2)_mNSO_2R^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C_1 - C_8 alkyl, C_2 - C_8 alkenyl, aryl, and C_1 - C_8 alkylaryl; and wherein R^6 and R^7 may independently combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may optionally have substituents selected from the group consisting of oxo, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, aryl, $-C_1$ - C_8 alkylaryl, $-C(O)C_1$ - C_8 alkyl, $-CO(O)C_1$ - C_8 alkyl, hydroxy, C_1 - C_8 alkoxy, $-C_1$ - C_8 alkylamine, amino, halo, and haloalkyl;

R^8 is hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_1 - C_8 alkylaryl, $-C(O)C_1$ - C_8 alkyl, or $-C(O)OC_1$ - C_8 alkyl; and wherein n is 0, 1, 2, 3 or 4 and m is 1, 2, or 3;

or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

2. The compound according to claim 1 wherein the A-ring is cyclohexyl.
3. A compound according to Claim 1 wherein the B-ring is selected from the group consisting of phenyl, pyridine, pyrimidine, pyrazine, and pyridazine.
4. A compound according to Claim 1 wherein the A-ring is piperidinyl.


5. A compound according to Claim 1 wherein E is an oxygen atom.
6. A compound according to Claim 1 wherein y is 0, 1, or 2, and R⁴ is independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, methoxy, ethoxy, methyl, ethyl, isopropyl, trifluoromethyl, trifluoromethoxy, phenyl, and benzyl.
7. A compound according to Claim 1 wherein z is 0, 1, or 2, and R⁵ is independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, methoxy, ethoxy, methyl, ethyl, isopropyl, trifluoromethyl, trifluoromethoxy, phenyl, and benzyl.
8. A compound according to Claim 1 wherein R¹ and R² are each independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl,




and wherein n is 1, 2, or 3.

9. The compound according to Claim 1 wherein R^6 and R^7 are each independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl:
10. A compound according to of Claim 1 wherein E is an oxygen atom, and R^6 and R^7 are both hydrogen atoms.
11. A compound according to Claim 1 wherein v is 1 or 2.
12. A compound according to Claim 1 wherein v is 1, m is 1, n is 1, y is 0 or 1 and z is 0 or 1.
13. A compound selected from the group consisting of:


6-{4-[2-(tetrahydro-pyran-4-yl)-ethylamino]-cyclohexyloxy}-nicotinamide,




6-[4-(3-Methyl-butylamino)-cyclohexyloxy]-nicotinamide,




6-[4-(2-Thiophen-2-yl-ethylamino)-cyclohexyloxy]-nicotinamide




4-[4-(3-Phenyl-propylamino)-cyclohexyloxy]-benzamide




Trans-6-(4-Benzylamino-cyclohexyloxy)-nicotinamide,




6-(1-Pyridin-2-ylmethyl-piperidin-4-yloxy)-nicotinamide




6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-nicotinamide




6-[1-(1H-Indol-2-ylmethyl)-piperidin-4-yloxy]-nicotinamide



4-(1-Benzyl-piperidin-4-yloxy)-benzamide,



4-[1-(3-Phenyl-propyl)-piperidin-4-yloxy]-benzamide



and a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer or a diastereomeric mixture thereof.

14. A compound according to Claim 1 wherein the pharmaceutically acceptable salt is the hydrochloric acid salt, the methanesulfonic acid salt, hydrobromide salt, the bisulfate salt or tartaric acid salt.

15. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to Claim 1 in association with a carrier, diluent and/or excipient.

16. A method for blocking a mu, kappa, delta or receptor combination (heterodimer) thereof in mammals comprising administering to a mammal requiring blocking of a mu, kappa, delta or receptor combination (heterodimer) thereof, a receptor blocking dose of a compound according to Claim 1, or a pharmaceutically acceptable salt, enantiomer, racemate, mixture of diastereomers, or solvate thereof.

17. A method of treating and/or preventing diseases related to obesity including irritable bowel syndrome, nausea, vomiting, obesity-related depression, obesity-related anxiety, smoking and alcohol addiction, sexual dysfunction, substance abuse, drug overdose, addictive behavior disorders, compulsive behaviors metabolic

diseases and symptoms thereof, and stroke, comprising administering a therapeutically effective amount of a compound of formula I.

18. A method of treating and/or preventing obesity and Related Diseases comprising administering a therapeutically effective amount of a compound of formula I to a patient in need thereof.

19. A method of suppressing appetite in a patient in need thereof, comprising administering a therapeutically effective amount of a compound of formula I.

20. A method of effecting weight loss in an obese patient comprising administering an effective amount of a compound of formula I or a pharmaceutically acceptable salt, solvate, racemate or enantiomer thereof.

21. A pharmaceutical composition for the treatment and/or amelioration of the symptoms associated with obesity and Related Diseases, containing as an active ingredient a compound of formula I according to Claim 1.

INTERNATIONAL SEARCH REPORT

International Application No
/US2004/038227

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C235/46 C07D211/46 C07D213/82 C07D401/12 C07D401/14
C07D405/12 C07D409/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	WO 03/082808 A (SUMITOMO PHARMACEUTICALS COMPANY, LIMITED; IMAZAKI, NAONORI; KITANO, M) 9 October 2003 (2003-10-09) abstract; claim 4; examples 25,160 -----	1-15
P,X	PATENT ABSTRACTS OF JAPAN vol. 2003, no. 12, 5 December 2003 (2003-12-05) & JP 2004 002367 A (TANABE SEIYAKU CO LTD), 8 January 2004 (2004-01-08) table 6; compound 10-8 -----	1-3,5-10
P,X	WO 2004/037800 A (GLAXO GROUP LIMITED; BEST, DESMOND, JOHN; BRUTON, GORDON; HEIGHTMAN, T) 6 May 2004 (2004-05-06) examples D36-D38 ----- -/--	1,3-7, 11,12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

18 April 2005

Date of mailing of the international search report

25/04/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hanisch, I

INTERNATIONAL SEARCH REPORT

International Application No

No. /US2004/038227

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99/33806 A (ORTHO-MCNEIL PHARMACEUTICAL, INC; CARSON, JOHN, R; CARMOSIN, RICHARD,) 8 July 1999 (1999-07-08) page 5, line 14 - line 18; claims -----	1-21
Y	EP 1 072 592 A (PFIZER INC; PFIZER LIMITED) 31 January 2001 (2001-01-31) cited in the application page 2, paragraph 1 - paragraph 2; claims 1,15,17 -----	1-21
Y	WO 01/74806 A (ASTRAZENECA AB; BROWN, WILLIAM; WALPOLE, CHRISTOPHER) 11 October 2001 (2001-10-11) page 2, line 7 - line 9; claims 1,8 -----	1-21
Y	WO 02/48122 A (ORTHO-MCNEIL PHARMACEUTICAL, INC) 20 June 2002 (2002-06-20) page 6, line 2 - line 11; claims -----	1-21
Y	WO 01/46174 A (ASTRAZENECA AB; BROWN, WILLIAM; WALPOLE, CHRISTOPHER) 28 June 2001 (2001-06-28) page 2, line 7 - page 5, line 3; claims 1-7; compound 6 -----	1-21
Y	WO 03/029233 A (RICHTER GEDEON VEGYESZETI GYAR RT; GALAMBOS, JANOS; NOGRADI, KATALIN;) 10 April 2003 (2003-04-10) page 2, line 3 - page 3, line 5; claim 1; example 4 -----	1-21
P,Y	WO 2004/026305 A (ELI LILLY AND COMPANY; BLANCO-PILLADO, MARIA-JESUS; CHAPPELL, MARK, DO) 1 April 2004 (2004-04-01) the whole document -----	1-21
P,Y	WO 2004/080968 A (ELI LILLY AND COMPANY; PEDREGAL-TERCERO, CONCEPCION; SIEGEL, MILES, GO) 23 September 2004 (2004-09-23) the whole document -----	1-21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2004/038227

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 16-20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
/US2004/038227

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03082808	A	09-10-2003	AU 2003220935 A1 EP 1500643 A1 WO 03082808 A1	13-10-2003 26-01-2005 09-10-2003
JP 2004002367	A	08-01-2004	NONE	
WO 2004037800	A	06-05-2004	AU 2003274053 A1 WO 2004037800 A1	13-05-2004 06-05-2004
WO 9933806	A	08-07-1999	AU 2009799 A CA 2316341 A1 EP 1049676 A1 JP 2001527068 T TW 476755 B WO 9933806 A1 ZA 9811842 A	19-07-1999 08-07-1999 08-11-2000 25-12-2001 21-02-2002 08-07-1999 23-06-2000
EP 1072592	A	31-01-2001	BR 0002515 A CA 2309434 A1 EP 1072592 A2 JP 2000344746 A JP 2004107324 A US 2002099216 A1	02-01-2001 28-11-2000 31-01-2001 12-12-2000 08-04-2004 25-07-2002
WO 0174806	A	11-10-2001	AT 289603 T AU 4498201 A BR 0109817 A CA 2403600 A1 CN 1432012 A ,C CZ 20023288 A3 DE 60109032 D1 EE 200200577 A EP 1286986 A1 HU 0300402 A2 JP 2003529598 T NO 20024777 A NZ 522143 A PL 357770 A1 WO 0174806 A1 SK 14242002 A3 US 2003199548 A1 ZA 200207973 A	15-03-2005 15-10-2001 21-01-2003 11-10-2001 23-07-2003 12-03-2003 31-03-2005 15-06-2004 05-03-2003 28-06-2003 07-10-2003 03-12-2002 25-06-2004 26-07-2004 11-10-2001 03-06-2003 23-10-2003 22-01-2004
WO 0248122	A	20-06-2002	AU 3401002 A WO 0248122 A2 US 2002123489 A1	24-06-2002 20-06-2002 05-09-2002
WO 0146174	A	28-06-2001	AT 275143 T AU 2418401 A BG 106805 A BR 0016591 A CA 2395300 A1 CZ 20022124 A3 DE 60013486 D1 EE 200200337 A EP 1242410 A1 ES 2225289 T3 HU 0300020 A2	15-09-2004 03-07-2001 31-03-2003 10-09-2002 28-06-2001 13-11-2002 07-10-2004 15-10-2003 25-09-2002 16-03-2005 28-05-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

US2004/038227

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0146174 A		JP 2003522741 T	29-07-2003
		MX PA02006063 A	06-02-2003
		NO 20022920 A	22-07-2002
		NZ 519986 A	25-06-2004
		PL 356780 A1	12-07-2004
		WO 0146174 A1	28-06-2001
		SI 1242410 T1	31-12-2004
		SK 8832002 A3	04-03-2003
		US 2003105110 A1	05-06-2003
		US 2004138225 A1	15-07-2004
		ZA 200204255 A	28-08-2003
WO 03029233 A	10-04-2003	HU 0103988 A2	28-05-2003
		EP 1438302 A1	21-07-2004
		WO 03029233 A1	10-04-2003
		JP 2005504126 T	10-02-2005
WO 2004026305 A	01-04-2004	AU 2003269980 A1	08-04-2004
		WO 2004026305 A1	01-04-2004
WO 2004080968 A	23-09-2004	WO 2004080968 A1	23-09-2004